

# MHD TIDAL WAVES ON A SPINNING MAGNETIC COMPACT STAR

YU-QING LOU<sup>1, 2</sup>

<sup>1</sup>National Astronomical Observatories, Chinese Academy of Sciences, A20, Datun Road, Beijing, 100012

<sup>2</sup>Department of Astronomy and Astrophysics, The Univ. of Chicago, 5640 S. Ellis Ave, Chicago, IL 60637  
 lou@oddjob.uchicago.edu

To appear in the *Astrophysical Journal Letters*

## ABSTRACT

In an X-ray binary system, the companion star feeds the compact neutron star with plasma materials via accretions. The spinning neutron star is likely covered with a thin “magnetized ocean” and may support *magnetohydrodynamic (MHD) tidal waves*. While modulating the thermal properties of the ocean, MHD tidal waves periodically shake the base of the stellar magnetosphere that traps energetic particles, including radiating relativistic electrons. For a radio pulsar, MHD tidal waves in the stellar surface layer may modulate radio emission processes and leave indelible signatures on timescales different from the spin period. Accretion activities are capable of exciting these waves but may also obstruct or obscure their detections meanwhile. Under fortuitous conditions, MHD tidal waves might be detectable and offer valuable means to probe properties of the underlying neutron star. Similar situations may also occur for a cataclysmic variable – an accretion binary system that contains a rotating magnetic white dwarf. This Letter presents the theory for MHD tidal waves in the magnetized ocean of a rotating degenerate star and emphasizes their potential diagnostics in X-ray and radio emissions.

*Subject headings:* MHD – stars: neutron – stars: oscillations – X-rays: stars – stars: binaries — stars: novae, cataclysmic variables

## 1. INTRODUCTION

For a magnetized spinning degenerate compact star covered with a thin plasma layer referred to as an *ocean*, various magnetohydrodynamic (MHD) tidal waves, excited by relevant processes, may exist. This scenario is plausible for an accretion binary system in which the Roche-overflow from a companion star accretes onto a spinning magnetized compact star. This Letter describes the theory of MHD tidal waves that propagate in a thin magnetized ocean of a spinning compact star. When combined with pertinent dynamic processes and emission mechanisms, such MHD tidal waves may leave indelible signatures of electromagnetic emissions in a wide spectrum ranging from the radio to GeV gamma rays and could provide key diagnostics that might offer clues and hints of otherwise complex phenomena. In general, MHD tidal waves may be excited gravitationally, thermally, electromagnetically, or by impacts of accreting materials. While accretion activities may stir up and sustain these waves, wave signals might be undesirably entangled with or overwhelmed by accretion noises. Once observationally established, these waves may be utilized to probe the structures in the *surface layers* of neutron stars and white dwarfs by extending or adapting the helioseismological techniques (Papaloizou & Pringle 1978; van Horn 1980; Saio 1982; Lou 2000, 2001b). Furthermore, damping properties of these waves might provide information of the stellar constituent materials and diffusion processes (Lou 1995).

## 2. THEORETICAL MODEL ANALYSIS AND RESULTS

MHD tidal waves may exist on spinning magnetic degenerate stars in either isolation or binaries (Chanmugam 1992; Phinney & Kulkarni 1994; Bildsten & Strohmayer 1999), although they might have a better chance to be de-

tected in high-mass and low-mass x-ray binaries (HMXBs and LMXBs) or cataclysmic variables. By strong gravity, the magnetized plasma ocean on a degenerate star of radius  $R$  is thin. As the ratio  $eR^3\Omega B/(GMm_p c)$  of electric force to surface gravity can be large for either neutron stars or white dwarfs in rotation (Goldreich & Julian 1969), where  $e$  is the electron charge,  $\Omega$  is the stellar angular spin rate,  $B$  is the poloidal magnetic field,  $c$  is the speed of light,  $G$  is the gravitational constant,  $M$  is the stellar mass and  $m_p$  is the proton mass, excess surface charges will be pulled upwards against gravity to form stellar magnetospheres. It is presumed that the ocean and an extended magnetosphere coexist on a compact star, that the magnetosphere maintains a pressure at the interface between the ocean and the magnetosphere much as the Earth’s atmosphere does on the ocean (Gill 1982), and that magnetic fields (idealized as a split monopole) are in and out radially across the thin ocean of thickness  $h$ . For a zonal magnetic field, the Alfvén-Rossby waves have been studied earlier in the solar/stellar context (Gilman 1969; Lou 1987, 2000).

For  $h/L \ll 1$  and a small Rossby number  $\mathcal{R} \equiv U/(2\Omega L)$  where  $U$  and  $L$  are typical horizontal velocity and spatial scales, the shallow-magnetofluid approach is valid. With a quasi-static vertical balance, the MHD tidal equations in spherical polar coordinates  $(r, \theta, \phi)$  are:

$$\frac{\partial v_\theta}{\partial t} - f v_\phi = -\frac{1}{R} \frac{\partial \Theta}{\partial \theta} - \frac{C_A^2}{c^2} \frac{\partial v_\theta}{\partial t}, \quad (1)$$

$$\frac{\partial v_\phi}{\partial t} + f v_\theta = -\frac{1}{R \sin \theta} \frac{\partial \Theta}{\partial \phi} - \frac{C_A^2}{c^2} \frac{\partial v_\phi}{\partial t}, \quad (2)$$

$$\frac{\partial \eta}{\partial t} + \frac{h}{R \sin \theta} \left[ \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{\partial v_\phi}{\partial \phi} \right] = 0, \quad (3)$$

$$\frac{\partial b_r}{\partial t} + \frac{B}{R \sin \theta} \left[ \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{\partial v_\phi}{\partial \phi} \right] = 0, \quad (4)$$

where variables  $v_\theta$ ,  $v_\phi$ ,  $\eta$  and  $b_r$  are the perturbations of  $\hat{\theta}$ -component velocity,  $\hat{\phi}$ -component velocity, ocean surface elevation, and  $\hat{r}$ -component of magnetic field, respectively,  $f \equiv 2\Omega \cos \theta$ ,  $g \equiv GM/R^2$ ,  $B$  and  $\rho$  are the Coriolis parameter at colatitude  $\theta$ , the surface gravity, the radial magnetic field, and the mean mass density, respectively,  $C_A \equiv B/(4\pi\rho)^{1/2}$  is the Alfvén speed,  $\Theta \equiv g\eta + Bb_r/(4\pi\rho)$  is proportional to the total pressure perturbation, and the *displacement current* effect is retained in case of  $C_A \gtrsim c$  (Lou 1995). The energy conservation follows from eqns. (1)-(4) with the energy flux density  $\vec{v}_\perp [\rho g\eta + Bb_r/(4\pi)]$  where  $\vec{v}_\perp \equiv (v_\theta, v_\phi)$ . By the  $\exp[i(\omega t + m\phi)]$  dependence with angular frequency  $\omega$  and integer  $m$ , the resulting equation of  $\Theta(\mu)$  becomes

$$\frac{d}{d\mu} \left( \frac{1 - \mu^2}{s^2 - \mu^2} \frac{d\Theta}{d\mu} \right) - \frac{1}{s^2 - \mu^2} \left( \frac{m s^2 + \mu^2}{s s^2 - \mu^2} + \frac{m^2}{1 - \mu^2} \right) \Theta + \frac{4R^2\Omega^2\Theta}{(gh + C_A^2)(1 + C_A^2/c^2)} = 0, \quad (5)$$

where  $\mu \equiv \cos \theta$  and  $s \equiv \omega(1 + C_A^2/c^2)/(2\Omega)$ . Variables  $v_\theta$ ,  $v_\phi$ ,  $\eta$  and  $b_r$  may be explicitly expressed in terms of  $\Theta(\mu)$ . The  $\Theta(\mu)$  solutions, regular at the two poles, involve the *Hough functions* (Hough 1897, 1898; Flattery 1967; Longuet-Higgins 1968; Chapman & Lindzen 1970). Given relevant physical parameters, all properties of MHD tidal waves can be derived computationally.

Without magnetic field, eqn (5) is equivalent to eqns (7) and (8) of Bildsten et al. (1996) with their two parameters  $q^2 \equiv s^{-2}$  and  $\lambda \equiv R^2\omega^2/(gh)$ . In their “traditional approximation”, the thin compressible layer is disturbed adiabatically, the radial component of the Coriolis force is ignored, the radial displacement is much less than the horizontal displacement, and  $\omega$  is much lower than the Brunt-Väisälä frequency. They focus on internal gravity modes of a rotating ocean. While internal magneto-gravity modes (Lou 1996) are excluded by invoking the quasi-static vertical balance, our incompressible formulation contains various MHD tidal waves in different regimes associated with the distinguished names of Kelvin (1879), Alfvén (1942), Rossby (1939), and Poincaré (1910), leading to the acronym of KARP waves.

For analytical simplicity, we hereafter invoke the  $\beta$ -plane approximation (Rossby et al. 1939) that retains the essential physics in the spherical geometry. In the local  $\beta$ -plane approximation of eqns (1) – (4) with  $\beta \equiv 2\Omega \sin \theta/R$ , a Cartesian coordinate system  $(x, y, z)$  is set up so that  $\hat{x}$ ,  $\hat{y}$  and  $\hat{z}$  point eastward (the sense of spin), northward and radially outward, respectively. One derives

$$\left[ \frac{\partial}{\partial t} \left( \frac{1}{c^{*2}} \frac{\partial^2}{\partial t^2} + \frac{f^{*2}}{c^{*2}} - \frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} \right) - \beta^* \frac{\partial}{\partial x} \right] v_\theta = 0, \quad (6)$$

where  $f^* \equiv f/(1 + C_A^2/c^2)$  is the *modified  $f$ -parameter*,  $\beta^* \equiv \beta/(1 + C_A^2/c^2)$  is the *modified  $\beta$ -parameter*, and  $c^{*2} \equiv (gh + C_A^2)/(1 + C_A^2/c^2)$  defines a characteristic speed  $c^*$ . For free MHD tidal waves of *small  $y$ -variation* scales and with  $v_\theta \propto \exp(i\omega t - ik_x x - ik_y y)$  where  $k_x$  and  $k_y$  are  $x$ - and  $y$ -component wavenumbers, the cubic dispersion relation is  $\omega^3 - [f^{*2} + c^{*2}(k_x^2 + k_y^2)]\omega - \beta^* c^{*2} k_x = 0$ . Global versions of KARP waves are contained in eqn (5).

### 3. EQUATORIALLY TRAPPED MHD TIDAL WAVES

By a latitudinal variation of  $f = \beta y$ , the *equatorial  $\beta$ -plane* becomes an effective *waveguide* for trapping, within the equatorial zone, MHD tidal waves that propagate in the  $\hat{\phi}$ -direction and this equatorial trapping effect becomes stronger in the regime of larger  $4R^2\Omega^2/[(gh + C_A^2)(1 + C_A^2/c^2)]$  in eqn (5) (Longuet-Higgins 1968). We examine main properties of such *equatorially trapped MHD tidal waves* below.

For *equatorially trapped Alfvén-Kelvin waves* (Thompson 1879; Alfvén 1942) around  $\theta = \pi/2$  that propagate only *eastward* at speed  $c^*$ , one must directly solve eqns (1) – (4) with  $v_\theta = 0$  to obtain

$$\{v_\phi, \eta, b_r\} = \{c^*, h, B\} \exp[-\beta^* y^2/(2c^{*2})] \mathcal{G}(x - c^* t), \quad (7)$$

where  $\mathcal{G}(\dots)$  is a *small arbitrary* dimensionless waveform and  $\beta^*$  appears in the Gaussian  $y$ -profile that decays at large  $|y|$  (e.g., Gill 1982).

With  $f = \beta y$  and a *nonzero*  $v_\theta \propto \exp(ik_x x - i\omega t)$  in the equatorial zone around  $\theta = \pi/2$ , eqn (6) becomes the Schrödinger equation of a harmonic oscillator. The normalized real solution takes the form of

$$v_\theta = \mathcal{N} \exp \left[ -\frac{\beta^* y^2}{2c^{*2}} \right] H_n[(\beta^*/c^*)^{1/2} y] \cos(k_x x - \omega t) \quad (8)$$

with  $H_n(\xi)$  being a Hermite polynomial of order  $n$  and the normalization constant  $\mathcal{N} \equiv 2^{-n/2}(n!)^{-1/2}[\beta^*/(\pi c^*)]^{1/4}$ . Variables  $v_\phi$ ,  $\eta$ , and  $b_r$  can be derived accordingly. The resulting *dimensionless* cubic dispersion relation for *equatorially trapped KARP waves* becomes

$$\tilde{\omega}^3 - (\tilde{k}_x^2 + n + 1/2)\tilde{\omega} - \tilde{k}_x/2 = 0, \quad (9)$$

where  $\tilde{\omega} \equiv \omega/(2\beta^* c^*)^{1/2}$ ,  $\tilde{k}_x \equiv k_x/(2\beta^* c^*)^{1/2}$ , and integer  $n \geq 0$  gives the number of nodes along  $y$ -direction. Results of eqn (9) are displayed in Figure 1 with the straight dashed line for the Alfvén-Kelvin mode (7) *added*. For  $n \geq 1$ , there are two classes of MHD tidal waves. The high-frequency class (upper solid lines) corresponds to *equatorially trapped Alfvén-Poincaré waves* (Poincaré 1910; Alfvén 1942) that propagate either eastward or westward but with *different* phase speeds  $\omega/k_x$  for a given  $|k_x|$ . The low-frequency class (lower solid lines) corresponds to *equatorially trapped Alfvén-Rossby waves* (Rossby et al. 1939; Alfvén 1942) with westward phase speeds; the group velocity  $d\omega/dk_x$  is eastward and westward for short- and long-waves, respectively. The  $n = 0$  case (dash-dot line) of eqn (9) corresponds to  $\tilde{\omega} = \tilde{k}_x/2 \pm (\tilde{k}_x^2/4 + 1/2)^{1/2}$  and is referred to as *equatorially trapped mixed Alfvén-Rossby-Poincaré waves* (Poincaré 1910; Rossby et al. 1939; Alfvén 1942) whose phase velocity can be to the east (plus sign) or west (minus sign) yet with a group velocity always to the east; for large  $k_x > 0$  it behaves more like Alfvén-Poincaré waves (plus sign), while for large  $|k_x|$  with  $k_x < 0$  it behaves more like Alfvén-Rossby waves (minus sign). There are no standing modes (i.e.,  $\tilde{k}_x = 0$ ) for equatorially trapped Alfvén-Kelvin and Alfvén-Rossby waves (Fig. 1). In contrast, for equatorially trapped Alfvén-Poincaré waves and mixed Alfvén-Rossby-Poincaré waves, *axisymmetric standing modes* exist with  $\tilde{k}_x = 0$  in eq. (9) and thus  $\tilde{\omega}^2 = n + 1/2$  where  $n = 0, 1, 2, \dots$ .

### 4. ESTIMATES AND DIAGNOSTICS

The preceding analysis is succinct with potential applications to diverse stellar settings. The estimates here

focus on typical neutron stars. One can extend the analysis to white dwarfs with different parameters. With  $M \sim 1.4M_\odot$  and  $R \sim 10^6\text{cm}$ , the surface gravity of a neutron star is  $g \sim 10^{14}\text{cm s}^{-2}$ . The ocean surface gravity wave speed is  $(gh)^{1/2} \sim 10^9 h_4^{1/2}\text{cm s}^{-1}$  and the Alfvén speed is  $C_A \sim 10^9 B_{12} \rho_5^{-1/2}\text{cm s}^{-1}$  for HMXBs or  $C_A \sim 10^6 B_9 \rho_5^{-1/2}\text{cm s}^{-1}$  for LMXBs, where  $h_\delta$ ,  $\rho_5$ , and  $B_\alpha$  stand for  $h$ ,  $\rho$ , and  $B$  in units of  $10^\delta\text{cm}$ ,  $10^5\text{g}$ , and  $10^\alpha\text{G}$ , respectively. For an extreme magnetar of  $C_A \sim 10^{12} B_{15} \rho_5^{-1/2}\text{cm s}^{-1}$ , relativistic and quantum effects are involved. For spin periods ranging from  $\sim 16\text{ms}$  to  $\sim 8.5\text{s}$ , the rotation speed  $\Omega R$  ranges from  $\sim 10$  to  $\sim 10^4\text{km s}^{-1}$ . It is clear that MHD tidal waves are important in the ocean dynamics of a spinning neutron star.

For a magnetar or neutron star in a HMXB, the situation of  $C_A \gtrsim c$ ,  $\Omega R$  and  $(gh)^{1/2}$  may happen with a stronger  $B$  and a less dense ocean. The inclusion of the displacement current term in eqns (1), (2), (5) and (6) is necessary (see the definitions of  $f^*$ ,  $\beta^*$ , and  $c^*$  following eq [6]) and the magnetic field is dynamically significant in MHD tidal waves besides potential diagnostic roles.

For a neutron star in a LMXB,  $C_A \ll c$ ,  $C_A < (gh)^{1/2}$ , and  $C_A \cong \Omega R$  unless  $B_9 \gg 1$ ,  $\rho_5 \ll 1$  and  $h_2 \ll 1$ . Thus,  $B$  affects only the rotational contribution to the speeds and frequencies of tidal waves significantly. Flow velocities of KARP waves are powerful to push magnetic fields around as  $v_\theta^2 + v_\phi^2 > b_r^2/(4\pi\rho)$ . Meanwhile, a  $B$  of  $\sim 10^8 - 10^9\text{G}$  would be strong enough to partially guide materials from an accretion disk onto the magnetic poles (White et al. 1983), leading to intense X-ray emissions as well as sporadic bursts (Bildsten & Strohmayer 1999).

KARP waves might be driven to excitation by radiative thermal or nuclear processes, by polar accretion impacts, by magnetic thrust and dragging, or by tidal interactions of the accretion disk, and in turn, are capable of periodically modulating thermal properties of the ocean and buffeting magnetic fields in polar regions to affect the height and cross section of polar accretions.

As an example of illustration, we now discuss possible diagnostics for equatorial MHD tidal waves of a neutron star in a LMXB. In spherical geometry,  $k_x = m/R$  for an integer  $m$  and  $\tilde{k}_x \sim 0.18m(M_{33}h_4)^{1/4}(R_6\nu_3)^{-1/2}$  for a small  $C_A$ , where  $\nu_3$ ,  $M_{33}$ , and  $R_6$  stand for  $\nu \equiv \Omega/(2\pi)$ ,  $M$ , and  $R$  in units of  $10^3\text{Hz}$ ,  $10^{33}\text{g}$ , and  $10^6\text{cm}$ , respectively. The Rossby radius of deformation is  $D_{\text{rossby}} \equiv (2\beta^*/c^*)^{-1/2} \sim 1.8 \times 10^5 (M_{33}h_4)^{1/4} \nu_3^{-1/2}\text{cm}$ . For  $R_6 \sim 1$ ,  $M_{33} \sim 2.8$  and  $\nu_3 \sim 0.5$ , equatorially trapped KARP waves influence a latitude range  $\gtrsim \pm 20^\circ$  across the equator.

Take a plausible geometry favorable for a detection. The spin and magnetic axes are misaligned with the two magnetic poles at latitudes outside  $\pm 45^\circ$ . The spin axis, likely aligned with the binary orbital axis, is taken to be perpendicular to the line of sight. During the phase of moderate and steady polar accretions from the surrounding disk, the stellar surface gives off a largely isotropic X-ray flux sufficiently away from the two magnetic polar hot spots. In this scenario, one may not directly detect either X-ray hot spots. *Instead, accretion-powered KARP waves trapped in the equatorial belt may give rise to relatively smooth large-scale ( $L \gtrsim R$ ) periodical modulations of X-ray brightness.* In principle, all these waves may coexist. Estimates are

given below for  $h_4 \sim 1$ ,  $R_6 \sim 1$ , and  $M_{33} \sim 2.8$ .

For  $m = 0$  or  $\tilde{k}_x = 0$ , dispersion relation (9) gives *standing* MHD tidal wave frequencies  $f_w$  as

$$f_w = \pm \frac{(2n+1)^{1/2}}{2\pi} \left[ \frac{2\Omega (GMh/R^2 + C_A^2)^{1/2}}{R (1 + C_A^2/c^2)^{3/2}} \right]^{1/2} \\ \cong \pm 5.1 \times 10^2 (2n+1)^{1/2} \nu_3^{1/2} (M_{33}h_4)^{1/4} R_6^{-1} \text{ Hz} \quad (10)$$

with  $n = 0, 1, 2, \dots$ . Given a detected  $f_w$ , the stellar spin rate of  $\nu$  can be inferred for  $n = 0$  – the special case of mixed Alfvén-Rossby-Poincaré standing waves (the dash-dot line in Fig. 1 with  $k_x = 0$ ). Or, for standing Alfvén-Poincaré waves of larger  $n \geq 1$ ,  $\nu$  would be smaller.

For nonaxisymmetric waves, the frequencies of equatorially trapped Alfvén-Kelvin waves (7) (the dashed straight line of Fig. 1) seen in an inertial reference frame are

$$f_w = 10^3 m \nu_3 + 1.3 \times 10^2 m (M_{33}h_4)^{1/2} R_6^{-2} \text{ Hz} \quad (11)$$

for a *small*  $C_A$ , where the corotation effect is included. With  $m = 1$  and a detected  $f_w$ , one can infer  $\nu$ . For  $m = 2$  and 3,  $f_w$  would be *precisely* 2 and 3 times the fundamental frequency.

For small  $\tilde{k}_x \sim 0.18m(M_{33}h_4)^{1/4}(R_6\nu_3)^{-1/2}$  with  $m \neq 0$ , the frequencies of equatorially trapped Alfvén-Rossby waves (lower solid lines of Fig. 1) as seen in an inertial reference frame are

$$f_w \cong 10^3 m \nu_3 - 1.3 \times 10^2 m (M_{33}h_4)^{1/2} R_6^{-2} / (2n+1). \quad (12)$$

With  $m = 1$ ,  $n = 1$ , and a detected  $f_w$ , one can infer  $\nu$ . For  $m = 2$  and 3,  $f_w$  would be *approximately* 2 and 3 times the fundamental frequency.

If only one apparent  $f_w$  were detected, it would be difficult to distinguish different KARP wave possibilities. A detection of two or more harmonics at 2 and 3 or more times the fundamental frequency may offer valuable hints. For nonaxisymmetric equatorially trapped Alfvén-Poincaré waves and mixed Alfvén-Rossby-Poincaré waves traveling around the globe, relation (10) must be modified to include the corotation effect  $m\nu$  and other  $m$ -dependence (Fig. 1); the overall  $m$ -dependence of these frequencies is not linear. Similarly, the *more accurate* version of relation (12) for Alfvén-Rossby waves does not contain a *strictly* linear  $m$ -dependence either (Fig. 1). By detecting more harmonics, one can constrain the type of KARP waves involved according to the dispersion relations (Fig. 1). As  $f_w$  given by eqns (10)–(12) vary with  $\nu$ , the expected stellar  $\nu$  spin-up in a LMXB (or unexpected variations, such as accreting X-ray pulsar Cen X-3) would lead to a  $f_w$  spin-up (or corresponding variations).

There have been recent theoretical development on stellar  $r$ -mode (Papaloizou & Pringle 1978; Saio 1982) instabilities caused by gravitational radiations (Andersson et al. 1999) to limit spin rates of hot young neutron stars at birth. The prospect that fast spinning neutron stars in LMXBs might emit gravitational waves (Bildsten 1999) observable by ground-based detectors currently under construction (e.g., the enhanced LIGO) offers another potential test for the KARP wave scenario advanced here by independent determinations of neutron star spin rates.

In the case that the stellar surface of a radio pulsar is covered with an ocean, various MHD tidal waves may modulate or interfere with radio pulse emissions from the magnetic polar regions by (quasi-)periodically varying polar

magnetic field strengths or changing magnetospheric conditions (Lou 2001a). This may lead to observable effects because magnetic and spin axes misalign for radio pulsars to shine. With  $B \gtrsim 10^{12}$  G for a pulsar,  $C_A$  and  $(gh)^{1/2}$  can be comparable and be both  $\gtrsim \Omega R$ . For equatorially trapped Alfvén-Poincaré waves, Alfvén-Kelvin waves, and the high-frequency branch of mixed Alfvén-Rossby-Poincaré waves, modulation timescales would be *shorter than or comparable to* radio pulse periods. For equatorially trapped Alfvén-Rossby waves and the low-frequency branch of mixed Alfvén-Rossby-Poincaré waves, modulation timescales would be *longer than* radio pulse periods. These are intriguing possibilities to be explored further.

This research was supported in part by grants from US NSF (AST-9731623) to the University of Chicago, by the ASCI Center for Astrophysical Thermonuclear Flashes at the University of Chicago under Department of Energy contract B341495, by the Visiting Scientist Programs at the Institute of Astronomy and Astrophysics, Academia Sinica (NSC-88-2816-M-001-0010-6) and at the National Center of Theoretical Sciences (Physics Division), National Tsinghua University, and by the Collaborative Research Fund from the NSF of China for Young Overseas Chinese Scholars (NSFC 10028306) at the National Astronomical Observatory, Chinese Academy of Sciences.

## REFERENCES

- Alfvén, H. 1942, *Nature*, 150, 405  
 Andersson, N., et al. 1999, *ApJ*, 516, 307  
 Bildsten, L. 1998, *ApJ*, 501, L89  
 Bildsten, L., & Strohmayer, T. 1999, *Phys. Today*, 52, No. 2, 40  
 Bildsten, L., Ushomirsky, G., & Cutler, C. 1996, *ApJ*, 460, 827  
 Chanmugam, G. 1992, *ARA&A*, 30, 143  
 Chapman, S., & Lindzen, R. S. 1970, *Atmospheric Tides, thermal and gravitational*. Gordon and Breach Science Pub., New York.  
 Flattery, T. W. 1967, *Hough Functions*. Tech. Rep. No. 21, Dept. of Geophys. Sci., The University of Chicago  
 Gill, A. E. 1982, *Atmosphere-Ocean Dynamics*. Academic Press  
 Gilman, P. A. 1969, *J. Atmos. Sci.*, 26, 1003  
 Goldreich, P., & Julian, W. H. 1969, *ApJ*, 157, 869  
 Hough, S. S. 1897, *Philos. Trans. R. Soc. London, Ser. A*, 189, 201  
 Hough, S. S. 1898, *Philos. Trans. R. Soc. London, Ser. A*, 191, 139  
 Laplace, P. S. 1778/1779. *Recherches sur plusieurs points du système du monde*. Mem. Acad. R. Sci. Paris [Reprinted in "Oeuvres," Vol. 9, pp. 71-183, 187-280, 283-310. Gauthier-Villars, Paris, 1893.]  
 Longuet-Higgins, M. S. 1968, *Philos. Trans. R. Soc. London, Ser. A*, 262, 511  
 Lou, Y. Q. 1987, *ApJ*, 322, 862  
 Lou, Y. Q. 1995, *MNRAS*, 275, L11  
 Lou, Y. Q. 1996, *Science*, 272, 521  
 Lou, Y. Q. 2000, *ApJ*, 540, 1102  
 Lou, Y. Q. 2001a, *ApJ*, 548, 460  
 Lou, Y. Q. 2001b, *ApJ*, 556, L121  
 Papaloizou, J., & Pringle, J. E. 1978, *MNRAS*, 182, 423  
 Pedlosky, J. 1987, *Geophysical Fluid Dynamics*. Springer-Verlag  
 Phinney, E. S., & Kulkarni, S. R. 1994, *ARA&A*, 32, 591  
 Poincaré, S. 1910, *Théorie des Marées, Leçons de Mécanique Celeste*, Vol. 3. Gauthiers-Villars, Paris  
 Rossby, C.-G., et al. 1939, *J. Marine Res.*, 2, 38  
 Saio, H. 1982, *ApJ*, 256, 717  
 Thompson, W. 1879, *Proc. Roy. Soc. Edinburgh*, 10, 92  
 van Horn, H. M. 1980, *ApJ*, 236, 899  
 White, N. E., Swank, J. H., & Holt, S. S. 1983, *ApJ*, 270, 711




FIG. 1.— Dispersion relations of KARP or MHD tidal waves.